


Name:
NHS No:
DoB:



Supported by



INDIVIDUAL HEALTH CARE PLAN FOR A CHILD OR YOUNG PERSON IN THE EDUCATION SETTING WHO HAS **TYPE 1** DIABETES



This care plan has been updated in December 2018 and is currently in line with National and International Guidelines.

CONTENTS:

1. CHILD/YOUNG PERSON'S INFORMATION	3
a. Child / Young Person Details.....	3
b. Family Contact Information	3
c. Essential Information Concerning This Child /Young Persons Health Needs	4
2. MONITORING GLUCOSE LEVELS	4
a. Continuous Glucose Monitoring (CGM).....	4
b. Flash Glucose Monitoring (FGS)	6
c. Blood Glucose Monitoring	6
3. INSULIN ADMINISTRATION WITH MEALS.....	7
4. HEALTHY EATING AND CARBOHYDRATE (CHO) COUNTING	7
5. INSULIN ADMINISTRATION.....	8
6. SUGGESTED DAILY ROUTINE.....	8
7. SPORTING ACTIVITY/ DAY TRIPS AND RESIDENTIAL VISITS.....	8
8. HYPOGLYCAEMIA	10
a. Treatment of Hypoglycaemia.....	11
9. HYPERGLYCAEMIA.....	12
a. Treatment of Hyperglycaemia For A Child/Young Person On Injections	12
b. Treatment of Hyperglycaemia for a Child/Young Person on Pump Therapy	13
c. Blood β -Ketone Testing Guide:	14
10. ADDITIONAL INFORMATION.....	14
11. CHECKLIST	15
12. TRAINING LOG.....	15
13. ADDITIONAL NOTES:.....	16
14. USEFUL RESOURCES	18
15. REFERENCES:	19

DEFINITIONS

IHCP	Individual Health Care Plan
CYP	Child or Young Person
HYPO	Hypoglycaemia
CHO	Carbohydrate
BG	Blood Glucose
CGM	Continuous Glucose Monitoring
FGS	Flash Glucose System

Name:
NHS No:
DoB:



This health care plan will capture the key information and actions that are required to support this child or young person (CYP) in school. Schools should make reasonable adjustments to facilitate prescribed medical care to allow a CYP with diabetes to participate in education on the same basis as their peers. This individual health care plan (IHCP) should be reviewed at least annually.

1. CHILD/YOUNG PERSON'S INFORMATION

a. Child / Young Person Details

Child's Name:				Year group:	
NHS number:				DoB:	
Nursery/School/College : Post code					
Child's Address: Town: County: Postcode					
Type of Diabetes:					
Other medical conditions:					
Allergies:					
Date:		Document to be Updated:			

b. Family Contact Information

Name		
Relationship		
Telephone Number	Home Work Mobile	
Email		
Name		
Relationship		
Telephone Number	Home Work Mobile	
Email		
Name		
Relationship		
Telephone Number	Home Work Mobile	
Email		

Name:
NHS No:
DoB:



c. Essential Information Concerning This Child /Young Persons Health Needs

Contacts			
	Name	Email	Contact Number
Children's Diabetes Nurses:			
Other Team Contact:			
Consultant Paediatrician:			
General Practitioner:			
Link Person in Education:			
Dietitian:			
Class Teacher:			
Health Visitor/School Nurse:			
SEN Co-ordinator:			
Other Relevant Teaching Staff:			
Other Relevant Professionals:			
Head teacher:			

This CYP has DIABETES, requiring treatment with (check which applies):

Multi-dose regime i.e. requires insulin with all meals:	<input type="checkbox"/>
Insulin Pump Therapy:	
3 injections a day (no injections in school):	<input type="checkbox"/>
2 injections a day (no injections in school):	
Other oral medication	
Other injectable medication	
Other medication:	

CYP with Diabetes will have to attend clinic appointments to review their condition. Appointments are typically every 3 months, but may be more frequent. These appointments may require a full day's absence. Education authority staff should be released to attend the necessary diabetes training sessions, in accordance with national guidance.

2. MONITORING BLOOD GLUCOSE (BG) LEVELS

BG monitoring is essential to achieving optimal diabetes management and must be familiar to school personnel. CYP with diabetes can monitor Glucose levels by the following ways; continuous glucose monitoring (CGM), flash glucose monitoring (FGS) and using a blood glucose monitor.

Monitoring glucose levels are an essential part of daily management; where ever possible the CYP should be encouraged to take responsibility for managing their own medicines and blood glucose (BG) equipment in school. They should be allowed to carry their equipment with them at all times and their equipment must not be shared.








a. Continuous Glucose Monitoring (CGM)

CGM is a small waterproof sensor that is inserted under the skin which measures glucose in the interstitial tissue every few minutes. The sensor sends the results to a transmitter. Unlike BG checking where you get a single measurement, with CGM, a graph is seen showing glucose levels with trends. Some sensors are linked to insulin pumps, some are stand-alone CGM. The trends give insight into the speed and direction the glucose is changing. Alarms can be set to alert.

**Examples of interpreting real-time CGM:
Medtronic (CGMS)**

Trend Arrow	What this means
↑ or ↓	Sensor Glucose has been rising or falling by about 1-2 mmol/L over the last 20 minutes
↑↑ or ↓↓	Sensor Glucose has rising or falling by about 2-3 mmol/L over the last 20 minutes
↑↑↑ or ↓↓↓	Sensor Glucose has been rising or falling by about 3 mmol/L over the last 20 minutes

Dexcom G4/G5/G6 (CGMS)






Trend Arrow G5 App	Trend Arrow G4/5 Receiver	What this means
	↑	Glucose rising 0.1-0.2mmol/L each minute or up to 2.5mmol/L in 15 minutes.
	↑↑	Glucose rapidly rising more than 0.2mmol/L each minute or more than 2.5mmol/L in 15 minutes.
	↘	Glucose is slowly falling 0.06-0.1 mmol/L each minute or up to 1.7mmol/L in 15 minutes.
	↓	Glucose is falling 0.1-0.2mmol/L each minute or up to 2.5mmol/L in 15 minutes.
	↓↓	Glucose is rapidly falling more than 0.2mmol/L each minute or more than 2.5mmol/L in 15 minutes.
	↗	Glucose slowly rising 0.06-0.1 mmol/L each minute or up to 1.7mmol/L in 15 minutes.
	→	Glucose is steady. Not increasing/ decreasing more than 0.06mmo/L per minute or up to 0.9mmol/L in 15 minutes.

b. Flash Glucose Monitoring (FGS)

This enables the glucose levels to be monitored by holding a handheld device (which is also a blood glucose monitor) over the sensor. This shows real-time glucose levels rather than BG and therefore there will always be a time lag in these glucose readings. Trend arrows show whether the levels are rising, decreasing or remaining stable. FGS do not continuously send glucose measurements to the reader.

Examples of interpreting:

Freestyle Libre Flash Glucose Monitor (FGS)

Trend Arrows	What this means
	Glucose is rising quickly (more than 0.1mmol/L per minute)
	Glucose is rising (between 0.06 and 0.1mmol/L per minute)
	Glucose is changing slowly (less than 0.06mmol/L per minute)
	Glucose is falling (between 0.06 and 0.1mmol/L per minute)
	Glucose is falling quickly (more than 0.1mmol/L per minute)
	The Glucose trend arrow may not always appear with your reading

c. Blood glucose monitoring

All CYP should have access to a blood glucose monitor whilst in the education setting. Some monitors may have a bolus advisor incorporated in the device. Some will check blood ketone levels as well, these are essential if on insulin pump therapy.

(Check which applies)

BG testing to be carried out by a trained adult, using a Fastclix / Multiclix device.	<input type="checkbox"/>
This child requires supervision with blood glucose monitoring.	<input type="checkbox"/>
This CYP is independent in BG testing.	<input type="checkbox"/>
This CYP may use CGMS	Please select
This CYP may use flash glucose monitoring	<input type="checkbox"/>
This CYP may use a meter with a bolus calculator	<input type="checkbox"/>

BG checking should be carried out:

- In class or if preferred, in a clean private area with hand washing facilities - **Not in a toilet.**
- Hands to be washed prior to the test.
- Blood glucose targets pre meal - mmol/L and - mmol/L 2 hours after meals.
- Blood glucose testing lancets and blood glucose strips should be disposed of safely.

There are a wide range of different blood glucose meters available, some have a built in automated bolus calculator, which will aid patients in determine CHO and correction doses.

3. INSULIN ADMINISTRATION WITH MEALS

Check if applies ☐ if not, go to section 5

Insulin can be delivered either by a pen device, or insulin pump therapy. The aim of the insulin pump is to mimic the pancreas as daily insulin requirements vary in each individual. They can either be attached to the wearer by a length of clear tubing (an infusion set) or as an adhesive patch pump attached to the skin by a small cannula.

The pump is set to deliver a continuous dose of fast-acting insulin over a 24 hour period (BASAL RATE); this basal rate will keep the BG stable when not eating. Extra fast insulin can be given via the pump, when the CYP is eating food and matched to the amount of carbohydrate that is eaten (BOLUS INSULIN). This is done by pushing buttons on the pump/handset (See page 14 additional information). Extra fast acting insulin can also be given as a CORRECTION DOSE if BG is above target range.

(Check which applies)

Insulin to be administered by a suitably trained adult, using a pen needle that complies with national and local sharps policy	<input type="checkbox"/>
Supervision is required during insulin injection administration	<input type="checkbox"/>
This young person is independent, and can self-administer the insulin injection	<input type="checkbox"/>
This CYP is on an insulin pump (see further information below and section 5 (page 8))	<input type="checkbox"/>

The CYP using either MDI or Insulin Pump therapy will need to count CHO and calculate insulin doses which ideally needs to be administered pre-meal.

(Check which applies)

They have a specific Insulin to carbohydrate (CHO) ratio (I:C)	<input type="checkbox"/>
They are on set doses of insulin	<input type="checkbox"/>

This procedure should be carried out:

- In class, or if preferred in a clean private area with hand washing facilities
- Should always use their own injection device; or sets.
- All used needles should be disposed of in accordance with the school's local policy

4. HEALTHY EATING AND CARBOHYDRATE (CHO) COUNTING

All CYP need a healthy and balanced diet for optimum growth and development. The meals eaten throughout a school day can make up a large proportion of some children's food intake for the whole day, perhaps including breakfast clubs, break times, lunch and after school clubs. As it's primarily the carbohydrate in food that raises the blood glucose level, it is therefore vitally important to count the carbohydrate accurately and administer the correct amount of insulin, the basis of good diabetes control. In some individuals the impact of dietary fat and protein should be considered in the calculation of the meal bolus dose.

A meeting can then be arranged with appropriate members of staff and parents to discuss who and how best supervision at 'meal times' can be implemented. In certain circumstances, it may be necessary for the dietitian to train staff in the basics of carbohydrate counting so they can support the child effectively.

The family will have received in-depth education regarding healthy eating and CHO counting from the diabetes team, and the CYP with diabetes should be allowed choice of whether to have a packed lunch or school meal. The process of CHO counting can be made easy utilising technology of their 'smart' phones.

Name:
NHS No:
DoB:



5. INSULIN ADMINISTRATION

Delivered via pen device: ☐ Delivered via insulin pump: ☐

Insulin Name	Time	Process
Please select		
Other:		
Insulin Name	Time	Process
Please select		
Other:		
Insulin Name	Time	Process
Please select		
Other:		
Insulin Name	Time	Process
Please select		
Other :		

NOTE: See 8 ☐

6. SUGGESTED DAILY ROUTINE

	Time	Note
Arrive School		
Morning Break		
Lunch		
Afternoon Break		
School finish		
Other		

Please refer to 'Home-school' communication diary ☐
Please refer to School planner ☐

7. SPORTING ACTIVITY/ DAY TRIPS AND RESIDENTIAL VISITS

Governing bodies should ensure that risk assessments, planning and arrangements are clear to ensure this CYP has the opportunity to participate in all sporting activities. School should ensure reasonable adjustments as required. Exercise will generally lower BG levels. Checking BG levels pre exercise is essential, and always consider what type of activity is being undertaken and what changes need to be made. For example:

Name:
NHS No:
DoB:



Before exercise:

BG Level		For this CYP:
4.0-4.9mmol/L	10-20 g carbohydrate regardless of type of exercise Start exercise once above 5.0mmol/l	
5.0-6.9mmol/L	10g carbs before aerobic Start anaerobic or interval exercise without snack	
7.0-15mmol/L	Start aerobic Start anaerobic or interval training but be aware blood sugars may rise.	
>15mmol/L	If hyperglycaemia is not explained e.g. not associated with recent meal check ketones* Consider 1/3 usual correction	

* If BG greater than 14 and/or Ketones above 0.6 mmol/l, follow advice on page 12/13 . Do not begin exercise until ketones less than 0.5 mmol/l.

Examples of Aerobic exercise in school: Football; Rugby; Hockey; Cross country

Examples of Anaerobic exercise in school: Sprinting; Gymnastics

After exercise:

Specific instructions for post exercise for this CYP:	
---	--

Check bloods immediately post exercise and before leaving school. Hypoglycaemia with exercise may occur at the time of the activity or may be delayed 7-11 hours later

Specific instructions for exercise for this CYP If on Insulin Pump therapy:	
---	--

If regularly having hypos in school after PE, school to discuss with family so changes can be made.

8. HYPOGLYCAEMIA

Hypoglycaemia ('Hypo') is best defined as a fall in BG level that exposes the CYP to potential harm and there may be no signs or numerical definition of hypoglycaemia for all CYP and situations.

Hypos are the commonest acute complication of Type 1 Diabetes. Blood glucose of below mmol/L requires urgent attention. It is important that all carers can recognise warning signs of hypoglycaemia and treat the low BG immediately and appropriately. A 'First Aid' hypo management pack must always be available.

Hypos can be symptomatic or asymptomatic, and are described as either mild or severe.

In rare circumstances an appropriate member of staff in school may be trained to administer the GLUCAGON injection. (resource 7)

Mild hypos events should be treated with oral glucose g glucose. The aim is to treat, and restore the BG level to above mmol/L.

Depending on the circumstances, this rapid acting glucose should be followed up by additional CHO to prevent reoccurrence of hypoglycaemia within 10-15 minutes.

Severe Hypos may result in loss of consciousness and / or a seizure. School personnel should have clear instructions for managing an episode of severe hypoglycaemia (see page 11) and use of rescue medication.



BG: Below 4 mmol/l.



INDIVIDUAL HYPO- SYMPTOMS FOR THIS CYP ARE:	Pale	<input type="checkbox"/>	Poor Concentration	<input type="checkbox"/>	Other:
	Sudden Change of personality	<input type="checkbox"/>	Sleepy	<input type="checkbox"/>	
	Crying	<input type="checkbox"/>	Shaking	<input type="checkbox"/>	
	Moody	<input type="checkbox"/>	CYP may say:		
	Hungry	<input type="checkbox"/>	Hypo unawareness		

How to treat a hypo:

- If possible, check BG to confirm hypo, and treat promptly: see 8a. If on FGS or CGMS.
- Do not send this child or young person out of class unaccompanied to treat a hypo.


A first aid management pack with fast acting glucose and long acting CHO should be kept in school containing fast acting glucose and long acting carbohydrate. Staff, and the CYP should be aware of where this is kept and it should be taken with them around the school premises; if leaving the school site; or in the event of a school emergency. It is the parent's responsibility to ensure this emergency box is adequately stocked; independent young people will carry hypo remedies with them.

Low BG levels can be caused by:

- Excess insulin
- Missed meal or Less CHO eaten than expected
- Exercise
- Excitement
- Hot weather
- Sleep
- Alcohol ingestion.

a. Treatment of Hypoglycaemia

BG BELOW MMOL/L



Additional information regarding hypoglycaemia for this CYP:

FOLLOWING RECOVERY CONSIDER WHAT HAS CAUSED THE HYPO

9. HYPERGLYCAEMIA
(High blood glucose)

Children and young people who have with diabetes may experience high blood glucose (hyperglycaemia) when the blood glucose levels are above **mmol/L**. The CYP may display the following symptoms:

Thirsty; frequent trips to the toilet; poor concentration; urinary accidents; mood changes

High BG levels can occur for the following reasons:


- Missed insulin dose
- Illness
- Extra CHO eaten
- Stress
- Hormones
- Injection site issues
- Less activity than normal

Please consider the cause before following advice from the flowcharts.

***** IF THIS CYP IS ON INSULIN PUMP THERAPY PLEASE REFER DIRECTLY TO 9b *****

If the child/young person is well, there is no need for them to be sent home, but parents/guardian should be informed at the end of the day that the child/young person has had symptoms of high blood glucose


a. Treatment of Hyperglycaemia For A Child/Young Person On Injections



Name:
NHS No:
DoB:

Additional information
regarding hyperglycaemia for
this CYP:

b. Treatment of Hyperglycaemia for a Child/Young Person on Pump Therapy



Name:
NHS No:
DoB:



c. Blood β -Ketone Testing Guide:

Below 0.6mmol/L

Between 0.6-1.5mmol/L

Above 1.5mmol/L

Normal range - low to 0.5mmol/L

Potential problems - SEEK ADVICE - 0.6 to 1.5mmol/L

High risk - SEEK URGENT ADVICE - 1.6mmol/L and above

10. ADDITIONAL INFORMATION

Additional information
regarding β Blood –Ketone
testing for this CYP:

- School to be kept informed of any changes in this child or young person's management.
- School exams or other assessment situations are associated with stress and increased risk of acute transient episodes of hypoglycaemia and hyperglycaemia that can affect performance. Specific arrangements may need to be in place for exams.
- The CYP with diabetes may be encouraged to wear identification stating they have diabetes. These are in the form of a bracelet, necklace, watch or medical alert card.
- There is no specific age at which a CYP with diabetes should be expected to take full responsibility for their diabetes in school.
- Specific extra support may be required for the CYP regarding educational, social and emotional needs- for example: during periods of instability: during exams: catching up with lessons after periods of absence: and counselling sessions.
- Storage of medication and supplies in school should be kept in line with local school policies. Teachers and school personnel have to be aware that the CYP with diabetes should have access to their devices and medication whenever needed.

Please use the box below for any additional information for this CYP, and document what is specifically important for him/her:

This IHCP has been initiated and updated in consultation with the CYP, family; diabetes specialist nurse and a member of staff from the educational setting.

Name:
NHS No:
DoB:



	Name	Signatures	Date
Young person			
Parents/Guardian			
Parents agreement to administration of medicine as documented on page 3 and 4			
Diabetes Nurse Specialist:			
School Representative:			
Health visitor/ School Nurse:			

11. CHECKLIST

First aid hypoglycaemia management pack	<input type="checkbox"/>	Insulin pen and appropriate pen needles	<input type="checkbox"/>
Glucose Gel (e.g. GlucoGel®/DextroGel®)	<input type="checkbox"/>	Cannula and reservoir for pump set change	<input type="checkbox"/>
Finger prick device, BG monitor and strips	<input type="checkbox"/>	Spare battery	<input type="checkbox"/>
Ketone testing monitor and strips	<input type="checkbox"/>	Up to date care plan	<input type="checkbox"/>
Snacks	<input type="checkbox"/>		<input type="checkbox"/>
Sharps bin	<input type="checkbox"/>		<input type="checkbox"/>

12. TRAINING LOG

Schools are responsible for adequately training their personnel about diabetes but the content of the training is the responsibility of the health care team and parents.

Staff Name	Training Delivered	Trainer	Date

Name:
NHS No:
DoB:



13. ADDITIONAL NOTES

Name:
NHS No:
DoB:



13. ADDITIONAL NOTES

14. USEFUL RESOURCES

- 1) **Carbs and Cals App:**
<https://www.carbsandcals.com/app/app>
- 2) **Carbs and Cals** book, by Chris Cheyette and Yello Balolia (Chello Publishing Ltd, ISBN no: 978-0-9564430-0-7) -
<https://www.carbsandcals.com/>
- 3) **Department of Education**
Supporting Pupils at School with Medical Conditions:
<https://www.gov.uk/government/publications/supporting-pupils-at-school-withmedical-conditions--3>
- 4) **Diabetes UK**
Diabetes in Schools Resources:
<https://www.diabetes.org.uk/Guide-to-diabetes/Schools/Diabetes-in-schoolsresources/>
Residential and Exams Packs:
<https://www.diabetes.org.uk/...to-diabetes/...diabetes/.../school-trip-and-exam-resource>
- 5) **DigiBete** - a place to help young people and families to manage Type 1 Diabetes
www.digibete.org
- 6) **Health and Safety (Sharp Instruments in Healthcare) Regulations 2013**
The relevant guidance can be accessed via the following link:
www.HSE.gov.uk/healthservices/index.htm
- 7) **How to Give an Injection of Glucagon (Glucogen) Video**
<https://www.digibete.org/video/give-injection-glucagon/>
- 8) **JDRF (Juvenile Diabetes Research Foundation)**
General Information Packs and Leaflets:
<https://jdrf.org.uk/living-with-type-1/information-packs-and-leaflets/>
Managing Type 1 diabetes effectively through University:
<http://uni.jdrf.org.uk/>
- 9) **National Children and Young People's Diabetes Network and JDRF E-learning module - Basic and Advanced Level**
<http://type1diabetestraining.co.uk/>
<https://jdrf.org.uk/for-professionals/school-pack/schools-e-learning-module/>
It is recommended that all staff complete the Basic Level. Staff who are carrying out blood glucose tests and giving insulin (either injections or via a pump) should also complete the Advanced Level.

15. REFERENCES:

An Individual Health Care Plan (IHCP) for a child or young person in an education setting who has diabetes within the Children and Young People's North West Diabetes Network (CYPNWDN) culminating in a national individual health care plan - M. Carson, D. Anderson, S. Singleton - ISPAD 2016: <http://medialibrary.ispad.cyim.com/mediatheque/media.aspx?mediald=17742&channel=9857>

Diabetes Guidelines for Schools, Colleges and Early Years Settings - East of England CYP Diabetes Network (versions 2013,2014,2018)

ISPAD Clinical Practice Consensus Guidelines 2018: <https://www.ispad.org/page/ISPADGuidelines2018>

Making every young person with diabetes matter. Department of Health, 2007

Managing Medicines in school and early Years Setting. Department of Health, 2005

Meeting Health Needs in Education and other Community Settings. RCN. January 2018

NICE clinical guideline NG18: Diabetes (type 1 and type 2) in children and young people, diagnosis and management, August 2015

Statutory Framework for Early Years Foundation Stage. Setting the standards for learning, development and care of children from birth to five.. Dept of Education, March 2017

Supporting pupils at school with medical conditions. Department of Education. September 2014
<https://www.gov.uk/government/publications/supporting-pupils-at-school-withmedical-conditions--3>

THIS CARE PLAN HAS BEEN REVIEWED AND REVISED IN 2018 BY:

Sandra Singleton, Children and Young People's Diabetes Specialist Nurse/ Team Leader
Dawn Anderson Children's and Young People's Diabetes Specialist Nurse
Margot Carson, Children and Young People's North West Diabetes Network Manager
Caroline McNicholas, Senior Diabetes Dietitian
Niccola Tate, Senior Diabetes Dietitian
Jonathan Maiden, Data Quality Manager, Children and Young People's North West Diabetes Network
Daniel Hyde IT Technical Support

ORIGINAL IHCP DESIGN BY A SUB-GROUP LEAD BY

Sandra Singleton, Children and Young People's Diabetes Specialist Nurse/ Team Leader with:
Margot Carson, Children and Young People's North West Diabetes Network Manager
Elaine McDonald, Children and Young People's Diabetes Specialist Nurse/ Team Leader
Dawn Anderson Children's and Young people's Diabetes Specialist Nurse
Linda Connellan, Children and Young People's Diabetes Specialist Nurse
Jill Cullen, Specialist Community Practitioner School Nursing
Jayne Johnson, Staff Nurse School Nursing.
Helen Nurse, Families with Diabetes National Network
Paula Maiden, Families with Diabetes National Network
Daniel Hyde, IT Technical Support

Acknowledgments:

The sub-group from the 'Children and Young People's North West Diabetes Network' would like to thank the following regional 'Children and Young People's Diabetes Network' teams for their helpful advice, support and input in the development and on-going review of this document:

Birmingham Children's Hospital (West Midlands)
Cambridge University Hospitals NHS Foundation Trust (East of England)
Chelsea & Westminster NHS Foundation Trust (London & South East Coast)
Glangwili General Hospital (Wales)
Leeds Children's Hospital (Yorkshire and the Humber)
North Somerset Community Partnership (South West)
North Tyneside General Hospital (North East and North Cumbria)
Nottingham Teaching Hospitals NHS Trust (East Midlands)
Oxford University Hospitals NHS Foundation Trust (Thames Valley)
Salisbury District Hospital (Wessex)



Review date: December 2019

Winner of the 'Excellence in Diabetes Specialist Nursing Award'
sponsored by:

SANOFI DIABETES 

at the Nursing Standard Nurse Awards 2015



Copyright © 2018 National Children and Young People's Diabetes Network